



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

143. Proposed by L. C. WALKER, A. M., Petaluma High School, Petaluma, Cal.

Find the area of greatest ellipse that can be inscribed in a given semicircle.

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa.; J. SCHEFLER, A. M., Hagerstown, Md.; and the PROPOSER.

Let the equation of the ellipse be

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \dots (1),$$

and of the circle, $x^2 + (y+b)^2 = r^2 \dots (2)$.

By eliminating x^2 from (1) and (2), then writing the condition for equal roots, we have

$$b^2 = a^2 - \frac{a^4}{r^2} \dots (3).$$

Required area $= \pi ab$, or say $u = a^2 b^2 = a^4 - (a^6/r^2)$. Hence for a maximum, we get

$$4a^3 = \frac{6a^5}{r^2}, \text{ or } a^2 = \frac{2r^2}{3}. \text{ Then the area} = \frac{2\sqrt{3}}{9} \pi r^2.$$

Also solved by JOSIAH H. DRUMMOND.

MECHANICS.

138. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, Gloucestershire, England.

A smooth elliptical tube is held in the vertical plane with its major axis inclined to the vertical. A particle is projected from the lowest point. Find the pressure on the tube at any point and the condition that the pressure may vanish at the highest point.

Solution by G. B. M. ZERR, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

The reaction on the tube is given by

$$v^2/\rho = R + X\rho(d^2x/ds^2) + Y\rho(d^2y/ds^2) = R + X(dy/ds) + Y(dx/ds).$$

Let the x -axis be vertical and let β be the inclination of the major axis to the vertical; then

$$(a^2 \sin^2 \beta + b^2 \cos^2 \beta)x^2 + (a^2 \cos^2 \beta + b^2 \sin^2 \beta)y^2 + 2(a^2 - b^2)xy \sin \beta \cos \beta = a^2 b^2$$

or $Ax^2 + By^2 + 2Cxy = a^2 b^2$, is the equation to the tube.

But $x = r \cos \theta$, $y = r \sin \theta$.

$$\therefore r = \frac{ab}{\sqrt{(A \cos^2 \theta + B \sin^2 \theta + 2C \sin \theta \cos \theta)}}$$